
Discover The Elegance Of A Hamiltonia



ASERETIC 10-25 Enalapril M leate-H drochlorothiazide dicated in patients

ett.

ntrandicated in patients frugs and in patients with Dosage must be inc. Evaluation of the hypertens For a Brief Sur-

avy Lot dizec ration: of f

action fed to previous treath and on fination is not let inite. all always include assessment of

enopy enal fue

USE IN PREGNANCY: When usinjury and even death to the develop the drechlorothiazide should be gnan. etus. the second and third timesters, ACL in actors of the actors of second second by the second sec

ABLETS ASERETIC® (ENALAPRIL MALEATE-HYDROCHLOROTHIAZIDE)

USE IN PREGNANCY: When used in pregnancy during the second and third trimesters, ACE inhibitors can cause injury and even death to the developing fetus. When pregnancy is detected, VASERETIC (Enalaprii Maleate-Hydrocthordinzide) should be discontinued as soon as possible. See WARNINCS, Fetal/Neonatal Morbidity and Mortality.

CONTRAINDICATIONS: VASERETIC is contraindicated in patients who are hypersensitive to any component of this product and in patients with a history of angioedema related to previous freatmr at with an angiotensin converting enzyme inhibitor. Because of the hydrochlorothiazide component, this product is contraindicated in patients with anuria or hypersensitive that the different date designed draw groups.

nent, mis product is continuated an patients in the too other sulfonamide-derived drugs.

WARNINGS: General; Enalapril Maleate; Hypotension: Excessive hypotension

WARNINGS: General: Enalapril Malichte: Hypotension: Excessive hypotension was rarely seen in uncomplicated hypertensive patients but is a possible consequence of enalapril use in severely salf-Volume depleted persons such as those treated vigorously with diuretics or patients on dialysis.

Syncope has been reported in 1.3 percent of patients receiving VASERETIC. In patients receiving enalapril alone, the incidence of syncope is 0.5 percent. The overall incidence of syncope may be reduced by proper titration of the individual components. (See PRECAUTIONS, Drug Interactions, and ADVERSE REACTIONS.)

Interactions, and ADVERSE REACTIONS.)

In patients with severe congestive heart failure, with or without associated renal insufficiency, excessive hypotension has been observed and may be associated with oliguria and/or progressive azotermia, and rarely with acute renal failure and/or death. Because of the potential fail in blood pressure in these patients, therapy should be started under very close medical supervision. Such patients should be followed closely for the first two weeks of treatment and whenever the dose of enalapril and/or diuretic is increased. Similar considerations may apply to patients with ischemic heart or cere-brovascular disease, in whom an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident. If hypotension occurs, the patient should be placed in the supine position and, if necessary, receive an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further doses, which

sient hypotensive response is not a contraindication to further doses, which usually can be given without difficulty once the blood pressure has increased after volume expansion.

after volume expansion.

Angioedema: Angioedema of the face, extremities, lips, tongue, glottis and/or larynx has been reported in patients treated with angiotensin converting enzyme inhibitors, including enalapril. This may occur at any time during treatment. In such cases VASERETIC should be promptly discontinued and appropriate therapy and monitoring should be provided until complete and appropriate therapy and monitoring should be provided until complete and sustained resolution of signs and symptoms has occurred. In instances where swelling has been confined to the face and lips the condition has generally resolved without treatment, although antihistamines have been useful in relieving symptoms. Angioedema associated with laryngeal edema may be fatal. Where there is involvement of the tongue, glottis or larynx, likely to cause airway obstruction, appropriate therapy, e.g., subcutaneous epinephrine solution 1:1000 (0.3 ml. to 0.5 ml.) and/or measures necessary to ensure a patent airway, should be promptly provided. (See ADVERSE REACTIONS.)
Patients with a history of angioedema unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see also CONTRAINDICATIONS).
Neutropenia/Agramulocytosis: Another angiotensin converting enzyme inhibitor, captopril, has been shown to cause agranulocytosis and bone marrow depression, rarely in uncomplicated patients but more frequently in patients with renal impairment especially if they also have a collagen vascular disease. Available data from clinical trials of enalapril are insufficient to show that enalapril does not cause agranulocytosis at similar rates. Marketing experience has revealed several cases of neutropenia or agranulocytosis in which a causal relationship to enalapril cannot be excluded. Penodic monitoring of white blood cell counts in patients with collagen vascular disease and renal disease, the patients with caution in severe renal disease. In patients with rouline in severe renal disease. Angioedema: Angioedema of the face, extremities, lips, tongue, glottis

Hydrochlorothiazide: Thiazides should be used with caution in severe renal disease. In patients with renal disease, thiazides may precipitate azotemia. Cumulative effects of the drug may develop in patients with impaired renal

truction.

Thiazides should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma.

Sensitivity reactions may occur in patients with or without a history of

Sensitivity reactions may occur in patients with or without a history of allergy or bronchial asthma.

The possibility of exacerbation or activation of systemic lupus erythematosus has been reported.

Lithium generally should not be given with thiazides (see PRECAU-TIONS, Drug Interactions, Enalagrii Maleate and Hydrochlorothiazide). Pregnancy, Enalagrii-Hydrochlorothiazide. There was no teratogenicity in rats given up to 90 mg/kg/day of enalagril (150 times the maximum human dose) in combination with 10 mg/kg/day of dyrochlorothiazide (2 ½ times the maximum human dose) or in mice given up to 30 mg/kg/day of enalagril (50 times the maximum human dose) in combination with 10 mg/kg/day of hydrochlorothiazide (2 ½ times the maximum human dose). At these doses, fetotoxicity expressed as a decrease in average fetal weight occurred in both species. No fetotoxicity occurred at lower doses; 30/10 mg/kg/day of enalagril-hydrochlorothiazide in rats and 10/10 mg/kg/day of enalagril-hydrochlorothiazide in rats and 10/10 mg/kg/day of enalagril-hydrochlorothiazide in rote.

When used in pregnancy during the second and third trimesters, ACE inhibitors can cause injury and even death to the developing fetus. When pregnancy is detected, VASERETIC should be discontinued as soon as possible. See Enalagril Maleate, Fetall/Neonatal Morbidity and Mortality. ACE inhibitors can cause fetal and neonatal morbidity and death when administered to pregnant women. Several dozen cases have been reported in the world literature. When pregnancy is detected, ACE inhibitors should be discontinued as soon as possible.

When pregnancy is detected, ACE infinitors should be uncommitted as a possible.

The use of ACE inhibitors during the second and third trimesters of pregnancy has been associated with fetal and neonatal injury, including hypotension, neonatal skull hypoplasia, anuria, reversible or irreversible renal failure, and death. Oligohydramnics has also been reported, presumably resulting from decreased fetal renal function; oligohydramnics in this setting has been associated with fetal limb contractures, craniofacial deformation, and hypoplastic lung development. Prematurity, intrauterine growth retardation, and patent ductus arteriosus have also been reported, although it is not clear whether these occurrences were due to the ACE-inhibitor exposure. These adverse effects do not appear to have resulted from intrauterine ACE-inhibitor exposure that has been limited to the first trimester. Mothers whose embryos and fetuses are exposed to ACE inhibitors only during the first trimester should be so informed. Nonetheless, when patients become pregnant, physicians should make every effort to discontinue the use of

Pregnant, physicians should make every effort to discontinue the use of VASERETIC as soon as possible.

Rarely (probably less often than once in every thousand pregnancies), no

*Registered trademark of Hospal Ltd.

10 mg

alternative to ACE inhibitors will be found. In these rare cases, the mothers should be apprised of the potential hazards to their fetuses, and serial ultrasound examinations should be performed to assess the intraamniotic envi-

ronment.

If oligohydramnios is observed, VASERETIC should be discontinued unless it is considered lifesaving for the mother. Contraction stress testing (CST), a non-stress test (NST), or biophysical profiling (BPP) may be appropriate, depending upon the week of pregnancy. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury.

the fetus has sustained irreversible finjury.

Infants with histories of in utero exposure to ACE inhibitors should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion or dailysis may be required as means of reversing hypotension and/or substituting for disordered renal function. Enalapril, which crosses the placenta, has been removed from neonatal circulation by peritoneal dialysis with some clinical benefit, and theoretically may be removed by exchange transfusion, although there is no experience with the latter procedure.

the latter procedure.

No teratogenic effects of enalapril were seen in studies of pregnant rats, and rabbits. On a mg/kg basis, the doses used were up to 333 times (in rats), and 50 times (in rabbits) the maximum recommended human dose. Hydrochlorothiazide; Teratogenic Effects: Reproduction studies in the rabbit, the mouse and the rat at doses up to 100 mg/kg/day (50 times the human dose) showed no evidence of external abnormalities of the fetus due to hydrochlorothiazide. Hydrochlorothiazide given in a two-litter study in rats at doses of 4 - 5.6 mg/kg/day (approximately 1 - 2 times the usual daily human dose) did not impair lertilify or produce birth abnormalities in the offspring. Thiazides cross the placental barner and appear in cord blood.

Nonteratogenic Effects: These may include fetal or neonatal jaundice, throm-boxtopenia, and possibly other adverse reactions which have occurred in

bocytopenia, and possibly other adverse reactions which have occurred in

the adult.

PRECAUTIONS: General: Enalapril Maleate: Impaired Renal Function: As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function may be anticipated in susceptible individuals. In patients with severe congestive heart failure whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, treatment with angiotensin converting enzyme inhibitors, including enalapril, may be associated with oliguria and/or progressive azotemia and rarely with acute renal failure and/or death. failure and /or death

failure and/or death.

In clinical studies in hypertensive patients with unilateral or bilateral renal artery stenosis, increases in blood urea nitrogen and serum creatinine were observed in 20 percent of patients. These increases were almost always reversible upon discontinuation of enalapril and/or diuretic therapy. In such patients renal function should be monitored during the first few weeks of

therapy.

Some patients with hypertension or heart failure with no apparent preexisting renal vascular disease have developed increases in blood urea and
serum creatinine, usually minor and transient, especially when enalapril has
been given concomitantly with a diuretic. This is more likely to occur in patients with pre-existing renal impairment. Dosage reduction of enalapril and/or discontinuation of the diuretic may be required.

Evaluation of the hypertensive patient should always include assess-

Hemodialysis Patients: Anaphylactoid reactions have been reported in

patients dialyzed with high-flux membranes (e.g., AN 69°) and treated con-comitantly with an ACE inhibitor. In these patients consideration should be given to using a different type of dialysis membrane or a different class of

inthippertensive agent.

Hyperkalemia: Elevated serum potassium (greater than 5.7 mEq/L) was Hyperkalemia: Elevaled serum potassium (graeter than 5.7 mdg/L) was observed in approximately one percent of hypertensive patients in clinical trials treated with enalapril alone. In most cases these were isolated values which resolved despite continued therapy, although hyperkalemia was cause of discontinuation of therapy in 0.28 percent of hypertensive patients. Hyperkalemia was less frequent (approximately 0.1 percent) in patients treated with enalapril plus hydrochlorothiazide. Risk factors for the development of hyperkalemia include renal insufficiency, diabetes mellitus, and the company of the properties of the development of hyperkalemia include renal insufficiency, diabetes mellitus, and the company of the development of hyperkalemia include renal insufficiency.

of hyperkalemia include renal insufficiency, diabetes mellitus, and the concomitant use of potassium-sparing duretics, potassium supplements and/or
potassium-containing salt substitutes, which should be used cautiously, if at
all, with enalaprii. (See Drug Interactions.)

Cough: Cough has been reported with the use of ACE inhibitors.

Characteristically, the cough is nonproductive, persistent and resolves after
discontinuation of therapy. ACE inhibitor-induced cough should be considered as part of the differential diagnosis of cough.

Surgery/Anesthesia: In patients undergoing major surgery or during anesthesia with agents that produce hypotension, enalapril may block
angiotensin II formation secondary to compensatory renin release. If
hypotension occurs and is considered to be due to this mechanism, it can
hypotension occurs and is considered to be due to this mechanism, it can
hypotension occurs and is considered to be due to this mechanism, it can
hypotension occurs and is considered to be due to this mechanism, it can
hypotension execution that of the performed at appropriate intervals.
All patients receiving thatacide therapy should be observed for clinical signs
of fluid or electrolyte imbalance hyponatremia, hypochloremic alkalosis, and
hypotalemia. Serum and urine electrolyte determinations are particularly
important when the patient is vomiting excessively or receiving parenteral important when the patient is vomiting excessively or receiving parenteral fluids. Warning signs or symptoms of fluid and electrolyte imbalance, irrespective of cause, include dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, confusion, seizures, muscle pains or cramps, muscular fatigue, hypotension, oliguria, tachycardia, and gastrointestinal distur-

cular latigue, hypotension, oliguna, tachycardia, and gastrointestinal disturbances such as nausea and vomiting. Hypokalemia may develop, especially with brisk diuresis, when severe cirrhosis is present, or after prolonged therapy. Interference with adequate oral electrolyte intake will also contribute to hypokalemia. Hypokalemia may cause cardiac arrhythmia and may also sensitize or exaggerate the response of the heart to the toxic effects of digitalis (e.g., increased ventricular urifability). Because enalapril reduces the production of aldosterone, concomitant therapy with enalapril attenuates the diuretic induced potassium loss (see Drug Interactions, Agents Increasing Serum Potassium).

Although any chloride deficit is generally mild and usually does not require specific treatment except under extraordinary circumstances (as in liver disease or renal disease), chloride replacement may be required in the

treatment of metabolic alkalosis

mg

Dilutional hyponatremia may occur in edematous patients in hot weathe appropriate therapy is water restriction, rather than administration of salt except in rare instances when the hyponatremia is life-threatening. In actual

except in rare instances when the hyponatrema is line-intreatening, in actual salt depletion, appropriate replacement is the therapy of choice. Hyperunicemia may occur or frank gout may be precipitated in certain patients receiving thiazide therapy. In diabetic patients dosage adjustments of insulin or oral hypoglycemic agents may be required. Hyperglycemia may occur with thiazide diuretics. Thus latent diabetes mellitus may become manifest during thiazide therapy. The antihypertensive effects of the drug may be enhanced in the postsympathectomic patient.

pathectomy patient.

If progressive renal impairment becomes evident consider withholding or discontinuing duretic therapy.

Thiazides have been shown to increase the urinary excretion of magne-

Thiazides have been shown to increase the urinary excretion of magnesium; this may result in hypomagnesemia. Thiazides may decrease urinary calcium excretion. Thiazides may decrease urinary calcium excretion. Thiazides may cause intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism. Marked hypercalcemia may be evidence of hidden hyperparathyroidism. Thiazides should be discontinued before carrying out tests for parathyroid function.

Increases in cholesterol and triglyceride levels may be associated with thiazide diuretic therapy.

Information for Patients: Angioedema: Angioedema, including laryngeal edema, may occur at any time during treatment with angiotensin converting enzyme inhibitors, including enalapril. Patients should be so advised and told to report immediately any signs or symptoms suggesting angioedema (swelling of face, extremities, eyes, lips, tongue, difficulty in swallowing or breathing) and to take no more drug until they have consulted with the prescribing physician.

physician.

Hypotension: Patients should be cautioned to report lightheadedness especially during the first few days of therapy. If actual syncope occurs, the patients should be told to discontinue the drug until they have consulted

patients should be told to discontinue the drug until they have consumed with the prescribing physician.

All patients should be cautioned that excessive perspiration and dehydration may lead to an excessive fall in blood pressure because of reduction in fluid volume. Other causes of volume depletion such as vomiting or diarrhea may also lead to a fall in blood pressure; patients should be advised to con-

Hyperkalemia: Patients should be told not to use salt substitutes containing

Hypertalemia: Patients should be told not to use salt substitutes containing potassium without consulting their physician.

Neutropenia: Patients should be told to report promptly any indication of infection (e.g., sore throat, fever) which may be a sign of neutropenia.

Pregunary: Female patients of childbearing age should be told about the consequences of second- and third-trimester exposure to ACE inhibitors, and they should also be told that these consequences do not appear to have resulted from intrauterine ACE-inhibitor exposure that has been limited to the first trimester. These patients should be asked to report pregnancies to their physicians as secon as receible.

their physicians as soon as possible.

NOTE: As with many other drugs, certain advice to patients being treated with VASERTIC is warranted. This information is intended to aid in the safe and effective use of this medication. It is not a disclosure of all possible adverse or intended effects

adverse or intended effects.
Drug Interactions: Enalapril Maleate; Hypotension—Patients on Diurctic Therapy;
Patients on diuretics and especially those in whom diuretic therapy was
recently instituted, may occasionally experience an excessive reduction of
blood pressure after initiation of therapy with enalapril. The possibility of
hypotensive effects with enalapril can be minimized by either discontinuing
the diuretic or increasing the salt intake prior to initiation of treatment with
enalapril. If it is necessary to continue the diuretic, provide medical supervision for at least two hours and until blood pressure has stabilized for at least
an additional hour. (See WARNINCS.)

Agents Causing Remin Release: The antihypertensive effect of enalapril is
augmented by antihypertensive agents that cause remin release (e.g., diuretics).

Other Cardiovascular Agents: Enalapril has been used concomitantly with beta adrenergic-blocking agents, methyldopa, nitrates, calcium-blocking agents, hydralazine and prazosin without evidence of clinically significant adverse interactions.

adverse interactions.

Agents Increasing Serum Potassium: Enalapril attenuates diuretic-induced potassium loss. Potassium-sparing diuretics (e.g., spironolactone, triamterene, or amiloride), potassium supplements, or potassium-containing salt substitutes may lead to significant increases in serum potassium. Therefore, if concomitant use of these agents is indicated because of demonstrated hypokalemia they should be used with caution and with frequent monitoring of serum potassium.

strated hypokalemia they should be used with caution and with frequent monitoring of serum potassium.
Lithium: Lithium toxicity has been reported in patients receiving lithium concomitantly with drugs which cause elimination of sodium, including ACE inhibitors. A few cases of lithium toxicity have been reported in patients receiving concomitant enalapril and lithium and were reversible upon discontinuation of both drugs. It is recommended that serum lithium levels be monitored frequently if enalapril is administered concomitantly with lithium.

Hutchchlorathyride: When administered concomitantly with lithium drugs. Hydrochlorothiazide; When administered concurrently the following drugs may interact with thiazide diuretics:

Alcohol, barbiturates, or narcotics—potentiation of orthostatic hypotension

may occur.

Antidiabetic drugs (oral agents and insulin)—dosage adjustment of the

Antidiabetic drugs (oral agents and insuini)—dosage adjustment of the antidiabetic drug may be required.

Other antihippertensive drugs—additive effect or potentiation.

Cholestyramine and colestipol resins—Absorption of hydrochlorothiazide is impaired in the presence of anionic exchange resins. Single doses of either cholestyramine or colestipol resins bind the hydrochlorothiazide and reduce its absorption from the gastrointestinal tract by up to 85 and 43 percent,

respectively.

Corticosteroids, ACTH—intensified electrolyte depletion, particularly


hypokalemia. Pressor amines (e.g., norepinephrine)—possible decreased response to presor amines but not sufficient to preclude their use. Skeletal muscle relaxants, nondepolarizing (e.g., tubocurarine)—possible to the production of the production

Skeletal muscle relaxants, nondepolarizing (e.g., tubocurarine)—possible increased responsiveness to the muscle relaxant. Lithium—should not generally be given with diuretics. Diuretic agents reduce the renal clearance of lithium and add a high risk of lithium toxcity. Refer to the package insert for lithium preparations before use of such preparations with VASERETIC.

Non-steroidal Anti-inflammatory Drugs—In some patients, the administration of a non-steroidal anti-inflammatory agent can reduce the diuretic, natruretic, and antihypertensive effects of loop, potassium-sparing and thiazide diuretics. Therefore, when VASERETIC and non-steroidal anti-inflammatory agents are used concomitantly, the patient should be observed closely to determine if the desired effect of the diuretic is obtained.

desired refer of undered sociations of Fertility: Enalapril in combination with hydrochlorothiazide was not mutagenic in the Ames microbial mutagen test with or without metabolic activation. Enalapril-hydrochlorothiazide did not produce DNA single strand breaks in an in vitro alkaline elution

APRIL 1994 • 160 • 4




Every cancer patient should have one.

Network (CAN) isn't. CAN is a nationwide program designed to help cancer patients of all ages travel free to or from recognized treatment. CAN arranges with corporations to utilize the empty seats on their corporate aircraft operating on business flights. This unique program allows cancer patients to travel in comfort and dignity. And best of all, the CAN program is free and it's easy. Since its inception in 1981, CAN has already helped thousands of patients across the country. If you'd like to find out more, call Corporate Angel Network at



PROGRAM PROFILE

University Plastic Surgeons/ Center For Reconstructive Surgery



icrovascular reconstruction has dramatically expanded the field of plastic surgery, opening doors for subspecialists who perform radical resections for the ablation of tumors or who attempt salvage of extremities that have been severely transmatized.

that have been severely cessfully promatized.

The Reconstructive Sur-

(b)

A 27-year-old who a sounced a conshot wound to the base of his thumbases left with (a) a composite defect anduling lost skin, muscle, andons need and bone; (b) simultaneous reconstruction of all these missing traduces and successfully performed as high a trade flap from his opposite upper arm manloying microsurgical technique.


ode o

age for

ger Team works closely with specialists in otolaryngology head and occasurgery, general surgery, orthopaedics, urologrehabilitation. Together they incorporate a varie blending innovative techniques that offer the grefunctional and aesthetic restoration, including

Hand Surgery

Re structive hand surgery is a modification potential techniques employed in several specialties; the impured the one of nerves and soft tissue, its restoucible community and hyaments. Surgical specialists perform the potation and reconstruction of complex has a commental hand deformities and hand hand hand hand.



 $A^{(SC)^{T}}$ n(vers) i Hospitai

USC Physician Access and Communication Exchange is sponsored by USC University Hospital, Childrens Hospital Los Angeles, The Doheny Eye Institute and USC/Norris Cancer Center. USC PACE is designed exclusively for use by physicians and their office staffs.

1.800.ASK.PACE

University of Southern California



We're advancing medicine.

A National Medical Enterprises Medical Center.

Reconstructive Microsurgery and Complex Wound Management

USC surgeons perform microsurgical free tissue transfer (using vascularized tissue flaps) to manage complex wounds which entails re-establishing circulation of the supplying arteries and draining veins between donor tissue and injury site.

Burn Reconstruction

Specialists in the secondary reconstruction of burn injuries use unique methods of restoring form and function. Surgeons have developed an operation that restores normal neck contour. and another that reconstructs ears. using materials which behave like living tissue. In facial reconstruction, they work

to restore balance to the patient's features, focusing not only on skin coverage but also on the structure of the facial skeleton to compensate for the loss of soft tissue and scar contracture.

Treatment of Craniofacial Deformities

The oromandibular region is particularly complex. Through advances in techniques and instrumentation, reconstructive surgeons can move the forehead, change the position of the eyes, realign the jaws and reshape the underlying structure to make dramatic differences in craniofacial deformities.

Reconstructive Breast Surgery

One alternative, the TRAM flap breast reconstruction procedure, involves the use of abdominal skin and muscles to form a new breast. Other options include autologous tissue replacement, implants, and tissue expanders followed by breast implant.

Post-Cancer Reconstruction

UNIVERSITY

PLASTIC SURGEONS

B. Chandrasekhar, M.D.

Bill Dougherty, M.D.

Susan E. Downey, M.D.

John Gross, M.D.

Larry S. Nichter, M.D.

John F. Reinisch, M.D.

Randolph Sherman, M.D.

Gerald M. Sloan, M.D.

Tad Wellisz, M.D.

extensive disfiguring tumor resection are grated implants for facial prosthetic reconstruction. In addition, musculoskeletal tumors in the upper and lower extremities that once required amputation may be treated through limb salvage procedures that

therapy, tumor resection, immediate implantation of bone allograft or endoprostheses, and microvascular coverage using free tissue transfer.

Close Communication With Referring Physicians

Members of the University Plastic Surgeons/ Center for Reconstructive Surgery are committed to maintaining close contact with referring physicians to ensure effective patient management for optimal recovery.

Patients who require benefitted by advances in microvascular techniques and osseointeincorporate antibiotic

School of Medicine

PHYSICIAN ACCESS AND COMMUNICATION EXCHANGE

APRIL 1994 • 160 • 4



CONTINUING MEDICAL EDUCATION CALENDAR

The Southern Medical Association is accredited by the Accreditation Council for Continuing Medical Education to sponsor continuing medical education for physicians.

APRIL 7-10 1994

FOCUS ON THE FEMALE PATIENT

LOCATION: Howey-in-the-Hills (Orlando), Florida

SPECIALTY: Primary Care, Ob/Gyn CME CREDIT: AMA Category I, 16 hours FEE: SMA Member \$310; Nonmember \$435

APRIL 15-17, 1994

ADDICTION MEDICINE

LOCATION: New Orleans, Louisiana

SPECIALTY: Primary Care, Psychiatry, Emergency

Medicine

CME CREDIT: AMA Category I, 15 hours FEE: SMA Member \$350; Nonmember \$475

APRIL 15-17, 1994

ORTHOPAEDIC UPDATE

LOCATION: Wrightsville Beach, NC SPECIALTY: Orthopaedic Surgery

CME CREDIT: AMA Category I, 14 hours FEE: SOA Member \$125; SMA Member \$225;

Nonmember \$350

APRIL 22-24, 1994

INFECTIOUS DISEASES LOCATION: Atlanta, Georgia SPECIALTY: Primary Care

CME CREDIT: AMA Category I, 15 hours FEE: SMA Member \$310; Nonmember \$435

MAY 20-22, 1994

CURRENT CONCEPTS IN ORTHOPAEDICS

LOCATION: Baltimore, Maryland SPECIALTY: Orthopaedic Surgery CME CREDIT: AMA Category I

FEE: SOA Member \$175; SMA Member \$275; Nonmember \$400; Resident or Fellow \$50;

Nonphysician \$75

JUNE 9-11, 1994

SOUTHERN ASSOCIATION FOR GERIATRIC MEDICINE, FOURTH ANNUAL MEETING

LOCATION: Hilton Head, South Carolina

SPECIALTY: Geriatrics

CME CREDIT: AMA Category I

FEE: To be announced

JUNE 12-16, 1994

EIGHTEENTH SYMPOSIUM ON LUNG DISEASE

LOCATION: Sea Island, Georgia

SPECIALTY: Pulmonary and Critical Care

Medicine

CME CREDIT: AMA Category I, 22 hours FEE: SMA/Southern ACCP Member \$415; Nonmember \$540; Nonphysician \$275

JUNE 16-18, 1994

MUSCULOSKELETAL IMAGING

LOCATION: Nevis, Caribbean

SPECIALTY: Radiology

CME CREDIT: AMA Category I, 17 hours FEE: SMA Member \$395; Nonmember \$520;

Nonphysician \$275

For more information, call Michelle Williamson, Southern Medical Association, 800-423-4992 or 205-945-1840—TODAY!

DECTION MALPRACTICE COVERAGE AT ITS BEST

- Effective and experienced management.
- An improved cash flow position immediately.
- \$1 million per occurrence/\$3 million aggregate per year.
- Affordable retroactive coverage.
- Remedial medical services designed to alleviate adverse medical/surgical results.

COMPARE AND SAVE

For further information, please call or write:



541 West Colorado Street, Glendale, California 91204 (818) 241-5119

Invest a stamp



Save a bundle

For the price of a stamp, you can get the latest edition of the federal government's free Consumer Information Catalog listing more than 200 free or low-cost government publications on topics such as federal benefits, jobs, health, housing, education, cars, and much more. Our booklets will help you save money, make money, and spend it a little more wisely.

So stamp out ignorance, and write today for the latest free Catalog. Send your name and address to:

Consumer Information Center Department SB Pueblo, Colorado 81009

A public service of this publication and the Consumer Information Center of the U.S. General Services Administration.



It's the professional edge in patient satisfaction and medicine compliance.

Prescribing the right medicine isn't enough. It's important to follow through and explain how and when to take it, precautions and side effects.

The National Council on Patient Information and Education (NCPIE) has free materials to help vou talk about prescriptions.

Write for free information on patient medicine counseling.



NCPLE 666 Eleventh Street, NW Suite 810 Washington, DC 20001

*AUGMENTIN**

amoxicillin/clavulanate potassium
See complete prescribing information in SmithKline Beecham Pharmaceuticals iterature or PDR. The following is a brief summary.
Indications and Usage: Augmentin sindicated in the treatment of infections caused by susceptible strams of the designated organisms in the conditions listed below.
Lower Respiration Tract Infectionscuaged by Blactarianses producing strains of Hemophilus influenzae and Moraxella (Branhamella) catarmals

Otitis Media caused by β -lactamase-producing strains of Hemophilus influenzae and

Moraxella (Branhamella) catarrhalis Sinusitiscaused by β-lactamase-producing strains of Hemophilus influenzaeand Moraxella

Blanhamella, catamhalis Skin and Skin Structure Intections caused by B-lactamase producing strains of Staphylo-cocus aureus E col and Klebsella spp Urnary Tract Infections caused by B-lactamase producing strains of E coli. Klebsiella spp

Unray Tract Infections caused by β-lactamase producing strains of £ coli, xieoseira syu and finerobacters spo. While Augmentin is indicated only for the conditions listed above, infections caused by ampcillin-susceptible organisms are also amenable to Augmentin treatment due to its amoscillin content. Therefore, insied infections caused by ampcillin-susceptible organisms and β-lactamase producing organisms susceptible to Augmentin should not require the addition of another antibotion. Bacteriological studies, to determine the causative organisms and their susceptibility to Augmentin should be performed together with any indicated surgical procedures. Therapy may be instituted prior to obtaining the results from bacteriological and susceptibility studies to determine the causative organisms and their susceptibility to Augmentin when there is reason to believe the infection may involve any of the β-lactamase producing organisms listed above. Once the results are known, therapy should be adjusted, if appropriate.

uyganisms issed above. Once the results are known, therapy should be adjusted, if appropriate
Contraindications: A history of allergic reactions to any penicillin is a contraindication
WARNINGS: SERIOUS AND OCCASIONALLY FATAL HYPERSENSITIVITY (ANAPHYLAC.
TOID). REACTIONS HAVE BEEN PEPORTED IN PATIENTS ON PROVICULIN THERAPY
ALTHOUGH ANAPHYLAXIS IS MORE FREQUENT FOLLOWING PARENTERAL THERAPY IN
HAS OCCURRED IN PATIENTS ON ORAL PROLICILINS THESE REACTIONS ARE MORE
LIKELY TO OCCUR IN INDIVIDUALS WITH A HISTORY OF PENICILLIN HYPERSENSITIVITY
AND/OR A HISTORY OF SENSITIVITY TO MULTIRE ALLERGENS. THERE HAVE BEEN
REPORTS OF INDIVIDUALS WITH A HISTORY OF PENICILLIN HYPERSENSITIVITY
AND/OR A HISTORY OF SENSITIVITY TO MULTIRE ALLERGENS. THERE HAVE BEEN
BEFORE INITIATING THERAPY WITH ANY PENICILLIN CARPICH, INDIVIDUALS SWITH A HISTORY OF PENICILLIN HYPERSENSITIVITY WHO
HAVE EXPERIENCED SEVERE REACTIONS WHEN TREATED WITH CEPHALOSPORINS
BEFORE INITIATING THERAPY WITH ANY PENICILLIN CARPICH INDIVIDIN'S SHOULD BE
MADE CONCERNING PREVIOUS HYPERSENSITIVITY PREACTIONS TO PENICILINS, CEPHAL
OSPORINS OR OTHER ALLERGENS AF AN ALLERGIC REACTION OCCURS, AUGMENT
SHOULD BE DISCONTINUED AND THE APPROPRIATE THERAPY INSTITUTED SERIOUS
ANAPPHYLACTION REACTIONS REQUIRE IMMEDIATE EMERGENCY TREATMENT
WITH EPINEPHRINE. DYGEN, INTRAVENOUS STEROIDS AND AIRWAY MANAGEMENT, INCLUDING INTUBATION, SHOULD ALSO BE ADMINISTERED AS
INDICATED.

PSEUDOMEROPHRINE DYGEN, INTRAVENOUS STEROIDS AND AIRWAY MANAGEMENT, INCLUDING INTUBATION, SHOULD ALSO BE ADMINISTERED AS
INDICATED.

PSEUDOMEROPHRINE DYGEN, INTRAVENOUS STEROIDS AND AIRWAY MANAGEMENT, INCLUDING INTUBATION, SHOULD ALSO BE ADMINISTERED AS
INDICATED.

INDICATE.

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including Augmentin, and has ranged in severity from mild to lifter-threatening. Therefore, it important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial

who present with draftines subsequent to the administration of antidacterial agents.

Treatment with antidacterial agents afters the normal flora of the colon and may permit overgrowth of clostrodia. Studies indicate that a toxin produced by Clostridium difficiers to primary cause of "artibiotic associated colins".

Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In who cases o pseudorinarious counts socially espond to up a social minder of another in moderate to severe cases, consideration should be given to management with fluids and electrolyses, protein supplementation and treatment with an antibacterial drug clinically effective against. C difficule colin.

Precautions: General: White Augmentin possesses the characteristic low toxicity of the

Procautions: General: While Augmentin possesses the characteristic low toxicity of the pencilial group of antibiotics pencidial group of admissions pencidial group of antibiotics pencidia assessment of organ system functions including renal hepatic and hematopoietic function, is advisable during prolonged therapy. A high perentage of patients with mononucleosis who received ampuchlin desire almost areas. Thus, ampound partial many functions with mononucleosis. The possibility of superinfections with mycotic or bacterial pathogens should be kept in mind during therapy. If superinfections occur lusually involving Pseudomonas or Candidal, the drug should be doscominued and/or appropriate therapy instituted. Drug Interactions: Pobenical decreases the renal fluxial's secretion of amoxicilin Concurrent use with Augmentin may result in increased and prolonged blood levels of amoxiciling.

amosciclin
The concurrent administration of allogurinol and ampicil\(^1\) in increases substantially the
incidence of rashes in patients receiving both drugs as compared to patients receiving
ampicilia alone it is not known whether this potentiation of ampicilin rashes is due to
allogurinol or the hyperuncema present in these patients. There are no data with
Augmentin and allogurinol administered concurrently
Augmentin should not be to-administered with Antabuse "disulfirami).

Augmentin should not be co-administered with Antabuse "Idisultram!
Carcinogenesis, Mutagenesis, Mutagenesis, Mutagenesis, Mutagenesis, Mutagenesis, Mutagenesis, magainesis, m

predictive of human response, this drug should be used during pregnancy only it clearly needed.

Labor and Delivery. Oral ampicullinic class antibiotics are generally poorly absorbed during labor. Studies in guinea pigs have shown that intravenous administration of ampicillini decreased the uterine tone, frequency of contractions, height of contractions and duration of contractions. Nowever, it is not known whether the use of Augmentinin humans during labor or delivery has immediate or delaved adverse effects on the fetus, prolongs the duration of labor or increases the likelihood that forceps delivery or other obstetrical intervention or resuscitation of the newborn will be necessary. Mussing Mothers: Ampicillin class antibiotics are excreted in the milk, therefore, caution should be exercised when Augmentin is administered to a nursing woman.

Adverse Reactions: Augmentin is generally well tolerated. The amporty of side effects observed in clinical trials were of a mild and transient nature and less than 3% of patients discontinued therapy because of thou related side effects. The most frequently reported aberse effects were damena/losse stools 19% in Russea (3% is sum lashes and uncarial 3% is committed the color and color and the stool and color and the color and color and the color and color and the color and the stool and the color and the stool and the color and the stool and the stool

tort, flatulence and headache
The following adverse reactions have been reported for ampicillin class antibiotics
<u>Gastrometernal</u> Diarrhea nausea vomiting indigestion gastritis stomatitis glossitis, black Thairy' tongue enterocolitis and pseudomembranous colitis Onset of pseudomembranous colitis symptoms may occur during or after antibiotic treatment (see WARNINGS)

pseudomembranous colints symptoms may occur during or after antibiotic treatment (see WARNINGS).

<u>Mypersensiturity reactions.</u> Shin rashes, urticaria, angioedema, serum sickness-like reactions furticate or shin rash accompanied by arthrustic, arthraligia, margiaja, and frequently fever! erythema multiforme (rarely Stevens Johnson Syndrome), and an occasional case of exfoliative dermatitis have been reported. These reactions ray be controlled with antihist ammers and if necessary, systemic corticosteroids. Whenever such reactions occur the drug should be discontinued, unless the opinion of the physician dictates otherwise. Serious and occasional fatal hypersensivity/anaphylactic reactions can occur with oral pencilin (see WARNINGS).


Legit A moderate rise in AST (SGOT) and/or ALT (SGPT) has been noted in patients treated with ampiculin class antibiotics but the significance of these frindings is unknown. Hepatic dystruction, including increases in serum transaminases (AST and/or ALT), serum bilirubin and or alkaline phosphatase; has been infequently reported with *Augmentin*. The histologic findings on liver biopsy have consisted of predominantly cholestatic, hepatocellular or mixed cholestatic hepatocellular changes. The onessel of significantion has occurred with time Hemic and Lymphatic Systems. Animal thomboxopolena, thromboxopolena comproblement action and agranulocytosis have been reported during therapy with pencillins. These reactions are usually reversible on decontinuation of therapy and are believed to be hypersensivity phenomena. A slight thromboxytosis was noted in less than be

believed to be hypersenstivity phenomena A signit thrombocytosis was noted in less than 1% of the patients treated with Augmentin Central Nerous System. Reversible hyperactivity, agitation, anwaty, insomnia, confusion, behavioral changes, and/or dizziness have been reported rarely BRS-AG L5A

STB SmrthKline Beecham Pharmaceuticals

When you suspect the presence of B-lactamaseproducing organisms in bronchitis and pneumonia*

The bugs stop here



Excellent climeat response rates in bronchitis and pneumonia

Latest in vitre insceptibility: still impressive H. influenzae. 100% and M. catarrhalis: 99.8%

Actively destro-B-lactamase

Ple see brief summar cost prindications, within on a fracent page.

Signith Kline Be

Classified Advertisements

The rate for each insertion is **\$8 per line** (six words per line) with five lines (**\$40**) minimum. Confidential box number charge: \$10 each month.

Classified display rates \$70 per inch. Maximum sizes: 1 column by 5 inches or 2 columns by 2 inches (2 1/8 inches per column). Larger classified ad space by special arrangements.

Copy for classified advertisements should be received not later than first of the month preceding issue. All copy must be typed or printed. • Classified advertisers using Box Numbers forbid the disclosure of their identity; your inquiries in writing will be forwarded to Box Number advertisers. Although The Western Journal of Medicine believes the classified advertisements in these columns to be from reputable sources, we do not investigate the offers made and assume no liability concerning them. The right is reserved to reject, withdraw, or modify all classified advertising copy in conformity with the decisions of the Advertising Committee.

Please Type or Print Advertising Copy

Classified Advertisements Are Payable in Advance

CLASSIFIED ADVERTISEMENTS
THE WESTERN JOURNAL OF MEDICINE
PO BOX 7602, SAN FRANCISCO, CA 94120-7602
(415) 882-3376
FAX: (415) 882-3379

PHYSICIANS WANTED

THE DIVISION OF FAMILY MEDICINE AT THE UNIVERSITY OF CALIFORNIA,

SAN DIEGO is seeking faculty members interested in combining clinical practice with clinical teaching. The positions will be at the Assistant Professor level, non-tenure track. The positions involve participating in a growing community-based, full spectrum Family Practice in a multicultural community with a full range of ages, socioeconomic, and health care payor systems. Opportunities for student and resident training are incorporated in the role. Candidates should be BC in Family Medicine with a strong commitment to teaching, clinical practice, and scholarship. Rank and salary, commensurate with qualifications and experience, based on UC scale. Interested candidates should send a CV and names of three references to:

UCSD
David Baughan, MD
Associate Clinical Professor
Division of Family Medicine (0807)
9500 Gilman Dr
La Jolla, CA 92093-0807

UCSD is an equal opportunity/affirmative action employer. All applications received by April 30 or thereafter until position is filled will receive thorough consideration.

NO HURRICANES, NO EARTHQUAKES! Just fresh air and happy living in north Idaho. An opportunity for a Primary Physician, Internist, Rheumatologist, Family Practitioner, or General Practitioner to take over an established practice at a minimal cost. Well equipped office, adjoining certified lab, experienced staff. New 170-bed hospital has state-of-the-art diagnostic and treatment facilities. Located in the foothills of the Rockies, Coeur d'Alene is a town of 25,000, a center of outdoor recreation, sports, and cultural activity. Please call Kelly at (208) 664-9234.

FAMILY PRACTICE. SALEM, OREGON. 20 plus multispecialty group in Oregon's beautiful Willamette Valley seeks BC/BE Family Practice physician. No OB. Good call rotation. Salary plus incentive, outstanding benefits. 45 minutes from Portland. Savor city life without the headaches. Natural wonders and recreational amenities abound. Contact Yvonne Carscadden at (503) 399-2470 or send CV to Salem Clinic, 2020 Capitol NE, Salem, OR 97303.

HAWAII. FAMILY PRACTITIONER needed for rural Oahu. Full-time position in nonprofit community health center. Desire dedication to work in multicultural setting. Obstetric skills preferred. Send letter and CV to Puanani Kalawa. Waianae Coast Comprehensive Health Center, 86-260 Farrington Highway, Waianae, HI 96792; (808) 696-7081.

SIMPLY THE BEST LOCUM TENENS PLACEMENT OPPORTUNITIES nationwide for Internal Medicine, Pediatrics, and Family Practice. Try a practice before you sign. Call The Locums' Company today at (800) 735-6286.

SOUTHERN CALIFORNIA. Great private practice opportunity in an affluent Orange county community. Primary Care practice with tremendous growth potential. BC in Family Practice or Emergency Medicine. Contact Dr. Joss, (714) 777-1773 or send CV to 21580 Yorba Linda Blvd, #102, Yorba Linda, CA 92687.

PHYSICIANS WANTED

DEAR DOCTOR:

If your goals include a quality lifestyle, a dynamic medical community, excellent schools/universities, and beautiful and affordable housing, then we have the ideal practice opportunity for you!

HOLY FAMILY HOSPITAL and the MEDICAL STAFF are working together to meet the healthcare demands of our 150,000 patient population. There is an IMMEDIATE need for BC/BE physicians in the following specialties:

- FAMILY PRACTICE
- INTERNAL MEDICINE
- GENERAL SURGERY
- PSYCHIATRY

SPOKANE, WASHINGTON (370K metro population) is the regional Healthcare and Cultural center offering a sound economy, unlimited outdoor recreation and a mild four season climate.

CONTACT with CV to: Nancy Chaffins



N. 5633 LIDGERWOOD SPOKANE, WA 99207 (509) 482-2133 FAX: (509) 482-2187

FAMILY PHYSICIAN-LEWISTON, IDAHO

Opportunity to join Family Practice group. Ambulatory and acute/inpatient care. No OB. Competitive income guarantee with opportunity to join full partnership. Send CV to Clearwater Medical Clinic, 1522 17th St, Lewiston, ID 83501, attn: Peggy Johnson; or call (208) 743-8416.

NORTHERN CALIFORNIA RECREATION AREA. Multispecialty group has immediate opening for an Internal Medicine practitioner with a particular interest in Critical Care. Clinical activities involve full range of Internal Medicine outpatient and inpatient practice. Beautiful northern California location offers abundant recreational opportunities as well as small town living. Competitive salary and comprehensive benefit package. Please send CV to 10978 Donner Pass Rd, Truckee, CA 96161.

SPOKANE, WASHINGTON OPPORTUNITY for BC/BE Family Practitioners and Pediatricians. Come enjoy this mild four season climate with unlimited year-round recreation. Take advantage of the excellent schools, universities, and affordable housing. Join this well established and growing multispecialty group offering choice practice opportunities within our satellite clinics. Advantages: favorable call schedules; an excellent benefits package and salary leading to early shareholder status; six weeks vacation and paid CME. Contact: Colleen Mooney; Rockwood Clinic; E 400 Fifth Ave, Spokane, WA 99202; (509) 838-2531, FAX (509) 455-5315.

(Continued from Page 390)

PHYSICIANS WANTED

NORTHWEST REGION

Physicians needed to join a multispecialty group, partnership, or solo practice, due to the explosive population growth. BC/BE physician specialties:

- FAMILY PRACTICE
- PEDIATRICS
- CARDIOLOGY
- PSYCHIATRY
- INTERNAL MEDICINE ORTHOPEDICS

Excellent practice opportunities offer quaranteed income and a strong referral base.

Contact with CV to:

Marge Becker, Owner/Manager **BECKER PHYSICIAN PLACEMENT** 4407 N Division, Ste 500 Spokane, WA 99207 (800) 945-4066/FAX (509) 484-6317

INTERNIST/FAMILY PRACTICE. Physician for well loved multidisciplinary Geriatric Health Center. Part of East San Francisco bay area community clinic network. Locum tenens position available. 60% time. Participation in hospital and call coverage desired. Resident teaching and UCSF clinical affiliation possible. Call (510) 644-6060.

CALIFORNIA-SAN LUIS OBISPO AREA. Position available in Urgent Care center on the beautiful central California coast. Competitive compensation, malpractice paid. Prefer physicians with experience or residency training in Emergency Medicine or Family Practice. Send CV to Medical Director, 283 Madonna Rd, Ste B, San Luis Obispo, CA 93405.

BC OR RECENTLY BE PHYSICIANS NEEDED. Los Angeles and Seattle metropolitan areas: Family Practice, Pediatrics, OB/GYN, Industrial Medicine, Urgent Care, and Internal Medicine. Northern Arizona: Invasive/Noninvasive Cardiologists and Internal Medicine. Call Chris Warner, Panther Enterprises Physician Recruiting and Placement, (800) 235-7601.

NEPHROLOGIST-BC/BE, for Internal Medicine/ Nephrology practice in San Francisco. Terms negotiable. Please send CV to A. Aronow, MD, 45 Castro St, San Francisco, CA 94114; (415) 565-6003.

ORTHOPEDIST-BC/BE, California licensed, needed for San Francisco based multispecialty medical practice. Primarily forensic medical practice (evaluation and expert witness evaluations in workers' compensation, personal injury, etc). Excellent opportunity for experienced clinician looking to emphasize evaluation and consultation practice as opposed to heavy treatment based practice. Full-time and parttime employment available, exceptionally high income potential, full benefits, and 401K. Submit CV and letter stating professional goals and availability to Administrator, 2040 Polk St, Box 337, San Francisco, CA 94109.

ROCKY MOUNTAIN WEST AND SOUTHWEST NEED PHYSICIANS. All specialties needed. Urban, rural, solo, group opportunities, all close to mountain recreation. Call Rita Longino at (800) 279-5267 or FAX CV to (800) 467-1246 or send CV to WHS, PO Box 2107, Corrales, NM 87048-2107.

(Continued on Page 392)





"I was worried I might be giving up continuity of patient care with an HMO. Not so with CIGNA. I have my own patient base, enabling me to give consistent, personalized medical care.'



"At CIGNA HealthCare, I have the freedom to enjoy an active social, as well as professional, life."

CIGNA HealthCare gives you the freedom to focus on what's most important...the practice of good medicine. We take care of the accounting, office management and more. And our locations offer you some of the best practice opportunities and lifestyles found anywhere. We currently have the following positions available:

SOUTHERN CALIFORNIA Los Angeles, Orange, Riverside & San Bernardino Counties

Dermatologist Family Physicians General Internists OB/GYNs Pediatricians

Urgent Care/FP & Peds

ARIZONA Phoenix & Tucson

Family Physicians General Internists Geriatric Internists Invasive Cardiologists OB/GYNs

Occupational Medicine Physicians Pediatricians Urgent Care Physicians

As part of our team, you'll enjoy a predictable schedule. What's more, CIGNA HealthCare offers competitive salaries, comprehensive benefits, and the resources of a healthcare leader. For more information, call or send your

California: CIGNA Medical Group, Physician Recruitment, 505 N. Brand Blvd., Suite 400-49, Glendale, CA 91203, (800) 468-9013.

Arizona: CIGNA HealthCare of AZ, Physician Recruitment. 11001 N. Black Canyon Hwy., Suite 401-49, Phoenix, AZ 85029, (800) 252-2471.



An Equal Opportunity Employer

PHYSICIANS WANTED

PHYSICIANS WANTED

COLORADO FAMILY PRACTICE PHYSICIAN

BC Family Physician to join a multisite community health center serving low income families in Northwest metropolitan Denver. Position includes faculty appointment at the University of Colorado Department of Family Medicine and resident training opportunities. Competitive salary and comprehensive fringe benefit package. Work in a teaching environment with an interdisciplinary team dedicated to promoting health in an underserved population. Please send CV to CLINICA **CAMPESINA FAMILY HEALTH SERVICES,** 1345 Plaza Ct N, Unit 1A, Lafayette, CO 80026. Or call Pete Leibig, Executive Director, at (303) 665-9310 for more information.

WESTERN MONTANA. BC/BE General Internist with interest in Primary Care wanted for active 50 plus physician multispecialty group. Unparalleled outdoor recreation and outstanding schools in university town of 45,000. Income guarantee, malpractice insurance, early partnership. Excellent fringe benefits. Position available now. Send CV to Administrator, Western Montana Clinic, PO Box 7609, Missoula. MT 59807.

TWO INVASIVE CARDIOLOGISTS AND FAMILY PHYSICIANS NEEDED. Unique opportunity. Gorgeous California central coast. Partnership within year. Send references/CV to Number 286, The Western Journal of Medicine, PO Box 7602, San Francisco, CA 94120-7602.

INTERNIST, BC/BE, to join four Internists in multispecialty practice in Portland suburb. Salary and benefits, with partnership available. Send CV to Beaverton Medical Clinic, Sterling Piepgrass, MD, 4690 SW Washington, Beaverton, OR 97229; (503) 644-3148.

FOR BEST RESULTS: CLASSIFIEDS. (415) 882-3376.

Classified Advertising is Affordable and Effective

❖

For details see the Classified Advertising form at the end of the journal. Or call

Classified Advertising: (415) 882-3376.

THE SOUTHWEST IS CALLING

Leading medical providers in Arizona, Nevada, Utah, Colorado, and New Mexico have asked us to find the best available BC/BE physicians in the specialties of Orthopedic Surgery, Internal Medicine, Family Practice, Pediatrics, Urgent Care, OB/GYN, Psychiatry, Hematology/ Oncology, Radiology, Occupational Medicine, ENT, MED/PED, Neurology, and Rheumatology. Top benefits, profit sharing, guarantee first two years, plus incentive pay, early shareholder. Please call CPR Tucson, (800) 658–9166, FAX (602) 322–2676, or in Phoenix, (800) 657–0354. FAX (602) 433–9548.

FAMILY PRACTICE, EMERGENCY MEDICINE, Internal Medicine, Locums, OB/GYN, Occupational Medicine, Pediatrics, and Urgent Care opportunities available in Minneapolis, St Paul, and leading communities within a 150-mile radius of the Twin Cities. Contact HealthSpan, Physician Services, 2810 57th Ave N, Minneapolis, MN 55430; (800) 248-4921 or (612) 574-7627.

SAN JOSE, CALIFORNIA. Busy academically affiliated hospital-based Occupational Medicine program with multidisciplinary focus is expanding practice in new clinic setting. Seeking full-time BC/BE experienced Occupational Medicine physician for advanced level clinical position. Competitive salary, excellent benefits. If interested, please send CV to Lynn E. Vail, MS, SCVMC, Division of Occupational Medicine, 750 S Bascom Ave, Ste 240, San Jose, CA 95128; (408) 299-8800, FAX (408) 993-1431.

PHYSICIANS WANTED



Health Care Recruiting NATIONWIDE Search Firm

James Ransom 160 NW Irving Ste 201-B PO Box 1221 Bend, OR 97709 (800) 323-1732 (503) 382-1732

SAN FRANCISCO BAY AREA. Energetic, recently trained BC/BE Pediatrician needed for 18 physician satellite of Palo Alto Medical Clinic, a 160 physician multispecialty group with a national reputation for innovation and excellence. Broad practice in growing bay area community. Excellent location with convenient access to San Francisco and Stanford. Competitive compensation/benefits package. Early partnership. Please send CV to David Hooper, PAMC, Fremont Center, 39500 Liberty St, Fremont, CA 94538

BOISE, IDAHO. IDAHO EMERGENCY PHYSICIANS is in the process of expanding our organization. We have two Urgent Care centers with plans to add an additional three centers by June 1994. We have full-time and part-time opportunities for experienced Primary Care physicians. Mid-level practitioners considered. Positions are salaried with superb benefits. Ownership potential. Ideal work week with flexibility. Extensive administrative support. Send résumé or CV for our return package. Recruiting Chairman, IEP, 2312 N Cole Rd, Ste B, Boise, ID 83704.

(Continued on Page 394)

To show you what we've been doing for the last 75 years, here are three perfect examples.



© 1993 National Easter Seal Society

Amanda, Cristy, and Brandy Jones, 10-year-old triplets, were born with varying degrees of disabilities, ranging from cerebral palsy, to motor, speech, and visual impairments. They were brought to Easter Seals for a pre-school program, and received a variety of occupational, physical and speech therapies. Helping people make the most of their abilities is what Easter Seals is all about. Which is why today, Amanda, Cristy and Brandy are just the three active fifth graders

Give Ability A Chance.

they should be.



FHP OFFERS PHYSICIANS THE BEST OF ALL WORLDS.

FHP Health Care is a highly successful managed care organization operating in prime locations in Southern California, Arizona, Nevada, Utah, New Mexico—even the tropical islands of Guam and Saipan. Whether you prefer big cities, beach communities, mountains, deserts, countrysides or the suburbs, you can have it with FHP.

We'll also provide what you want in the world of your practice. FHP's Primary Care Delivery System enables you to make the key decisions that affect your practice. You'll have all the support you need, from physician specialists to ancillary professionals. And you'll share in the rewards of a successful practice.

You'll also have three weeks paid vacation, fully paid malpractice insurance, CME, excellent compensation and comprehensive benefits including retirement programs, 401k and more.

Giving Physicians More Of What They Want.



With a 30 year history of solid growth. FHP ranks among America's most successful HMOs. For information about opportunities in Southern California, call (800)241-4347; in Utah, (800)283-8884; in Arizona, (800)535-4347; in New Mexico, (800)444-4347; or in Nevada, Guam or Saipan, (800)336-2255. If you are unable to call, please send your resume to our Corporate Office: FHP Health Care, Professional Staffing Dept., 9900 Talbert Ave., Fountain Valley, CA 92708. Equal Opportunity Employer.



PART OF AN OPERATION THAT'LL MAKE YOU FEEL BETTER

As an Air Force Reserve physician, you'll experience all the rewards of providing care. And then some. Because as part of our nation's vital defense team, you'll help protect the strength and pride of America. In the Air Force Reserve, you'll feel the excitement a change of pace brings as you gain the prestige of military rank and the privilege of working with some of the world's best medical professionals—in a program that's flexible enough to fit your schedule. The Air Force Reserve. It's a great way to serve.

Call: (800) 253-6189



(Continued from Page 392)

PHYSICIANS WANTED

Western States OPENINGS

Many multispecialty groups and hospitals have asked us to recruit for over 300 positions of various specialties. Both permanent and locum tenens. Send CV to Western States Physician Services, 5627 E. Kings Canyon, Ste 156, Fresno, CA 93727, or call (800) 873-0786.

EXCELLENT OPPORTUNITY FOR INTERNAL MEDI-CINE PHYSICIAN in private practice multispecialty physician group in San Francisco. Income guarantee. No investment. Forward CV to Box 263, The Western Journal of Medicine, PO Box 7602, San Francisco, CA 94120-7602.

CALIFORNIA MULTISPECIALTY. Urgent Care Director, Dermatologist, Radiologist, Otorhinolaryngologist, General Surgeon, Cardiologist, Internal Medicine, Pediatrician, Gastroenterologist, Orthopedist, General/Family Practitioner, Obstetrician/ Gynecologist. Excellent opportunity for physicians to join 400 member multispecialty medical group serving the greater Los Angeles, San Fernando Valley, and Orange county areas. Over 40 locations. Large fee-forservice and prepaid practice, no Medi-Cal. Excellent compensation program based on guarantee plus incentive, profit sharing and pension plan. Group provides health, dental, life, and malpractice. Partnership in real estate and medical corporation available. Send CV to Ron McDaniel, Assistant Administrator, Mullikin Medical Center-5, 17821 S Pioneer Blvd, Artesia, CA

PHYSICIANS WANTED

IDAHO

Does having free time, three and four days off in a row, no on-call, a flexible schedule, and a competitive compensation package sound good to you? How about living in beautiful Idaho and experiencing a great life style? We are currently seeking full-time Family Practice physicians to staff our Family Practice/Minor Emergency clinics throughout Idaho. Call Lora Meadows at (800) 688–5008 or send CV and inquiries to Primary Health, 727 E River Park Ln, Boise, ID 83706.

PUGET SOUND. BC/BE Family Practitioners to affiliate with 100 plus physician multispecialty group, located 25 miles north of Seattle. Both walk-in clinic and traditional Family Practice position available. Opportunity for exceptional personal and professional lifestyle. Positions offer competitive salary with excellent benefits. Available immediately. Send CV to J. G. Finley, MD, The Everett Clinic, 3901 Hoyt Ave, Everett, WA 98201.

PHYSICIANS. We are a six site Community Health Center located in central California. We have an opening at our clinic near Bakersfield for an Internist and a Family Practitioner. Competitive salary, revenue from hospitalization, excellent benefits, daytime hours, and limited call. Loan repayment available through NHSC. Contact George Laven, MD or George Johnston at (805) 845–3731.

PHYSICIANS WANTED

EMERGENCY PHYSICIAN

CODY, WYOMING has a year round position available for an Emergency Physician beginning September of 1994. Work in the beautiful Rocky Mountain region, located only 50 miles from Yellowstone Park. Area abundant in fishing, camping, boating, and all outdoor sports. NO STATE INCOME TAX!! For more information call (307) 578-2488 or send CV to Gary Bishop, Administrator, West Park Hospital, 707 Sheridan Ave, Cody, WY 82414.

ENT/HNS-UTAH. Small college town two hours from Alta and Snowbird. Nearby mountain biking, hunting, and desert exploring. Excellent 84-bed hospital with supportive medical staff. Well equipped office. Write to David Hilding, MD, Ste 1, Hospital Dr, Price, UT 84501.

HEAT WAVE? HAWAII, CALIFORNIA, UTAH, South Carolina, Florida, and Arizona clinics need Family Practitioners, Internists, Noninvasive Cardiologists and OBs. Hawaii—Hematologist/Oncologist, Gastroenterologist, \$130,000. Benefits. Clear speech. CVs to LAM Associates, Dept W, 444 Hobron Ln, #207H, Honolulu, HI 96815-1229; Pat Lam, (800) 258-4526, (808) 947-9815, 11 am-8 pm PST.

CALIFORNIA. MONTEREY BAY. Full-/part-time positions available with Monterey Bay's largest and most successful Urgent Care network. Generous guarantee, incentive plan, and benefit package. Malpractice covered. Practice in California's most beautiful coastal recreational area. BC/BE Emergency Medicine or Family Practice specialists preferred. Contact Rob Colyn, VP of Personnel, Doctors on Duty Medical Clinics, 19000 Portola Dr, Ste 100, Salinas, CA 93908; (408) 455-2424.

ASSOCIATE IN PEDIATRICS. Kern Medical Center, Bakersfield, California, a teaching hospital affiliated with UCLA and UCI Schools of Medicine, seeks an Associate in the Division of Pediatrics. Prerequisites include eligibility or certification by the American Board of Pediatrics, strong interest in teaching and qualifications for faculty appointment in UCLA Department of Pediatrics. Comprehensive salary and benefit package. A part-time private practice is permitted. Medical center is in central California, a mid-sized urban community with moderate cost of living. Send CV and inquiries to Navin Amin, MD, Chairman, Department of Family Practice/Pediatrics, Kern Medical Center, 1830 Flower St, Bakersfield, CA 93305.

Ads Get Results!

Call Classified

Advertising
for more information.

(415) 882-3376

PHYSICIANS WANTED

MARKS MEDICAL PLACEMENTS Joyce Marks-President

Immediate opportunities for BC/BE physicians in South Carolina, Kentucky, Kansas, Georgia, Texas, Nevada, Louisiana, Washington, Southern, and Northern California; the BEST locations. Family Practice, Internal Medicine, OB/GYN, Pediatrics, Dermatology, Physician Assistants. Nurse Practitioners

Please send CV to:

Joyce Marks-President Marks Medical Placements 2170 Century Park East, #806 Los Angeles, CA 90067

Or Call: (310) 556-2126, FAX (310) 556-1604.

AT NO COST TO PHYSICIAN

FAMILY PRACTICE RESIDENCY, FULL-TIME FAC-ULTY POSITION. University of Wyoming, Family Practice Residency Program-Casper. This outstanding rural Family Practice residency program seeks a physician scholar to join our faculty. The program has nine faculty members, 24 residents, and a competent support staff. The residency is affiliated with the Wyoming Medical Center, the referral hospital for the state, and the program is housed in an outstanding, freestanding facility. Responsibilities include resident supervision in ambulatory and inpatient settings, participation in scholarly activities faculty development, and direct patient care. Requirements include BC in Family Practice, ability to obtain Family Practice credentials including Obstetrics, and a strong desire to teach rural Family Practice. Completion of a teaching fellowship in Family Practice and experience with Area Health Education Centers are highly desirable. Applications should include a current CV and three references. Academic rank in based on the individual's experience, and a competitive salary and benefits are offered. The community of Casper and the state of Wyoming can provide an enjoyable life-style which is unequaled elsewhere. The University of Wyoming is an equal opportunity/affirmative action employer. Please contact Joe K. Schoeber, MD, Program Director, University of Wyoming, Family Practice Residency Program-Casper, 1522 East "A" St, Casper, WY 82601; (307) 266-3076.

ENJOY THE GREAT SOUTHWEST. Thomas-Davis Medical Centers, one of Arizona's fastest growing multispecialty groups is seeking BC/BE Internal Medicine, Family Practice, OB/GYN, Hematology/Oncology, and Urgent Care Physicians for our Tucson location. 200 physicians strong, TDMC has provided health care services in Arizona for more than 75 years. For more information regarding this practice opportunity, contact William Trainor, TDMC, 6135 N 7th St, Phoenix, AZ 85014, or call (800) 539-2529, or FAX CV to (602) 530-6255.

BC/BE FAMILY PRACTITIONER with interest in OB and a commitment to caring for the underserved needed to join clinics in Grandview and Walla Walla, Washington, and Hermiston, Oregon. All clinics located in rural settings with abundant recreational activities. Competitive salary, Public Health Service loan repayment slots, and excellent benefit package including paid malpractice. Contact Mayling Chin, Recruiting Specialist, or Ann Garza, Director of Personnel at Yakima Valley Farm Workers Clinic, PO Box 190, Toppenish, WA 98948, or (509) 865-5898.

LOOKING FOR THE PERFECT POSITION? Place a classified advertisement under "POSITION WANTED" and get noticed. Classified advertising is a cost-effective way to reach 46,000 physicians nation-wide. Call (415) 882-3376 to place your ad.

PHYSICIANS WANTED

CALIFORNIA

Primary Care Physicians and Radiologists needed to work as locum tenens statewide. High salary, paid malpractice. Work whenever and wherever you wish. Permanent placements as well. INTERIM Physicians/Western Physicians Registry; Northern California, Jim Ellis, Director, (800) 437-7676. Southern California, Tracy Zweig, Director, (800) 635-3175.

IDAHO-NONINVASIVE CARDIOLOGIST. 18 physician, multispecialty clinic with attached 44-bed hospital, seeks noninvasive Cardiologist to join busy practice in a growing community. Facility is physician owned, partnership available after two years, above average compensation. Outstanding recreational opportunities. Close to Sun Valley and wilderness areas. Please send CV to Brent Bodily, PO Box 1233, Twin Falls, ID 83301; (208) 733-3700.

CALIFORNIA. Ambulatory Care/Family Practice-CEP is a partnership of Ambulatory Care and Emergency Medicine specialists. We currently staff nine free-standing clinics in northern, central, and southern California, and have upcoming openings for physicians specializing in Family Practice. We offer shared ownership, excellent compensation, health, and disability insurance, retirement program, and many other benefits. For more information, send CV or call John Gravette, CEP, 2101 Webster St, Ste 1050, Oakland, CA 94612; (800) 842-2619.

COASTAL CALIFORNIA-SAN LUIS OBISPO. Family Practice and Internal Medicine, BC/BE. Join our busy Primary Care group practice. Located midway between Los Angeles and San Francisco on the central coast of California. San Luis Obispo offers a variety of cultural/recreational activities, a beautiful coastline, vineyards, and a local university. Call or send CV to Fran Coughlin, RN, MS, Ambulatory Care Services, San Luis Obispo General Hospital, Box 8113, San Luis Obispo, CA 93403-8113. Phone (805) 781-4936, FAX (805) 781-1096.

GREAT FAMILY PRACTICE OPPORTUNITY in small town in the north San Francisco bay area. Practicemoderately paced, broad range Family Practice group. No medicaid. Competent and caring staff with very little turnover. Stable patient base. Affiliated with residency program. Excellent opportunity for growth. Call 1:4 with inpatient optional. Well equipped office with view of bay, next to state park and par course. Base salary plus productivity bonus. Willing to provide housing. One year contract with option for partnership. Community-best of both worlds; small town life within 30 minutes of San Francisco. Affordable housing, good schools, safe neighborhood. Four seasons climate with mild winter. Outstanding recreational and cultural amenities. Boating, windsurfing, biking, and fine restaurants. Requirements-BC/BE Primary Care, minimum one year contract. To start August 1, 1994. Please contact Walt Zaks, Family Health Center, 1440 Military West, Benicia, CA 94510 or phone (707) 745-0711.



PHYSICIANS WANTED

THE VA MEDICAL CENTER, PRESCOTT, ARIZONA

is seeking a BC/BE Internist or Family Practice Physician. Regular hours. Experience with Holter monitors, treadmill testing, and/or endoscopy desirable but not necessary. Prescott is a lovely area with beautiful surroundings, clean air, pleasant lifestyle, wonderful dry four season climate. Interested physicians may call (602) 445-4860, ext 6640, or send CV to Chief, Medical Service, VA Medical Center, Prescott, AZ 86313.

EOE

WYOMING INTERNIST NEEDED. Step into full practice with established Internist with guaranteed salary first year, buy-in second year. Practice real Internal Medicine like the old days. Great small town life but close enough to a major metropolitan area. No state income tax. Incredible outdoor activities with camping, flyfishing, world-class skiing nearby. Send reply to Box 289, Western Journal of Medicine, PO Box 7602, San Francisco, CA 94120-7602.

PHYSICIANS. The California Department of Corrections has positions available for Internal Medicine, Primary Care, General Practice, and Family Practice at the California Correctional Center at Susanville in beautiful Lassen County in Northern California. For more information, contact Stephen O'Barr, (916) 257-2181, ext 1213, or FAX your résumé to (916) 322-5864. EOE.

FAMILY PRACTICE. BC/BE PHYSICIANS sought by well established, physician-guided managed health care provider in Arizona. Primary Care environment. Phoenix and Tucson area opportunities. Shared call, predictable hours, attractive salary, outstanding benefits, and incentive programs. For more information, call (800) 535-4347 or send CV to FHP Health Care, Professional Staffing, PO Box 52078, Phoenix, AZ 85072-2078. FHP may also have clinical and management opportunities available in California, Utah, New Mexico, Nevada, and Guam. An equal opportunity employer M/F/D/V.

EMERGENCY MEDICINE UNIVERSITY POSITION.

The University of California, Davis, School of Medicine, is recruiting for a full-time faculty. The position will be at the Assistant or Associate Professor level. The Division of Emergency Medicine and Clinical Toxicology is undergoing rapid academic development. A residency training program in Emergency Medicine was begun in 1990 and currently has 24 residents. The UCDMC Emergency Department provides comprehensive emergency service and is a major trauma center in northern California. The Department is a Paramedic Base Station and Training Center, and in addition, has an active helicopter service and regional poison center. Candidates must be BC/BE in Emergency Medicine and be eligible for licensure in California. A letter outlining interests and experience in addition to a CV and the names of five references should be sent to Robert W. Derlet, MD, Chair, Emergency Medicine Search Committee, FOLB I, University of California, Davis, Medical Center, 2315 Stockton Blvd, Sacramento, CA 95817. Opened until filled. No applications will be accepted after April 30, 1994. The University of California is an Affirmative Action/Equal Opportunity Employer.

BOOKS

We have two new FREE CATALOGS.

- 1. Medical Books Catalog
- 2. Allied Health and Nursing Books Catalog

Call, write or come in for your free copy today.

(206) 543-6582 • Fax 685-3921

University Book Store

301 South Campus Center University of Washington Seattle, WA 98195 Open Mon.-Fri. 8 AM-5 PM; Thurs. nights 'til 8; Sat. 10 AM-4 PM

SERVICES

BIMONTHLY MEDICAL NEWSLETTER On HOW TO PASS BOARD OF INTERNAL MEDICINE

For a **FREE** first copy mail your address label (self-sticking) with four (29 cents) stamps to

Medical Newsletter

389 E Mt Pleasant Ave Livingston, New Jersey 07039.

Also available-similar newsletter on "How to Pass Board of Geriatric."

NEW LAWS PUT DOCTORS AT RISK

Do you understand the many new federal laws and regulations which apply to medical offices? We will perform an on-site survey of your office and practice, find the potential problems, and help you correct them. Our fees are lots easier to take than a million dollar lawsuit or large fine.

COMPLIANCE SURVEY SERVICE (800) 605-9535

MEDICAL LITERATURE RESEARCH. Physician/educator/editor will do customized literature searching and analysis for medical or legal problems. Call Bill Milburn, MD, (303) 823-5083.

SUPPORT STAFF

CALIFORNIA: NURSE PRACTITIONERS AND PHYSICIAN ASSISTANTS PROVIDED for your clinical setting. Locum tenens and permanent placements through Western Practitioner Resources, California's most active, most respected placement service for Nurse Practitioners and Physician Assistants. For further information contact Heidi Bourne. RN. Director. (800) 345-5859.

MEDICAL EQUIPMENT

GOING OUT OF BUSINESS SALE. Serono-Baker 9000 CBC Analyzer (2 years old), \$8,000. Titmus Vision Tester (4 months old), \$1,100. DuPont Coumtrak \$250. Call (801) 753-7525.

ADVERTISERS:

Is today the deadline?

FAX us your ad!

Our FAX number is:

(415) 882-3379

"...IF ALL WOMEN **OVR 50** HAD REGULAR MAMMOGRAMS, THEIR DEATH RATE FROM BREAST CANCER WOULD DROP BY A THIRD."

-ACOG

Take the first step. Call 1-800-ACS-2345 for information that can help you make an impact.

— give the word.— MAMMOGRAM Reference: 1, Jones PH, et al. Once-daily pravastatin in patients with primary hypercholesterolemia: a dose-response study. Clin Cardiol. 1991;14:146-151.

PRAMACHOL® (Pravastatin Sodium Tablets)

CONTRAINDICATIONS

Hypersensitivity to any component of this medication.

Active liver disease or unexplained, persistent elevations in liver function tests (see WARNINGS).

Pregnancy and lactation. Atherosclerosis is a chronic process and discontinuation of lipid-lowering drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hypercholesterolemia. Cholesterol and other products of cholesterol biosynthesis are essential components for fetal development (including synthesis of steroids and cell membranes). Since HMG-COA reductase inhibitors decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol, they may cause fetal harm when administered to pregnant women. Therefore, HMG-CoA reductase inhibitors are containdicated during pregnancy and in nursing mothers. Pravastatin should be administered to women of childbearing age only when such patients are highly unlikely to conceive and have been informed of the potential hazards. If the patient becomes pregnant while taking this class of drug, therapy should be discontinued and the patient apprised of the potential hazard to the fetus.

WARNINGS

patient apprised of the potential hazard to the tetus.

WARNINGS

Liver Enzymes: HMG-CoA reductase inhibitors, like some other lipid-lowering therapies, have been associated with biochemical abnormalities of liver function. Increases of serum transaminase (ALT, AST) values to more than 3 times the upper limit of normal occurring on 2 or more front necessarily sequentially ocasions have been reported in 1.3% of patients treated with pravastatin in the U.S. over an average period of 18 months. These abnormalities were not associated with cholestasis and did not appear to be related to treatment duration. In those patients in whom these abnormalities were believed to be related to pravastatin and who were discontinued from therapy, the transaminase levels usually fell slowly to pretreatment levels. These biochemical findings are usually asymptomatical though worldwide experience indicates that anorexia, weakness, and/or abnormal pain may also be present in rare patients.

when the set benefit with praesisted of the related to resident of the related to the comment of the comment of

Other Drugs: During clinical trials, no noticeable drug interactions were reported when PRAVACHOL was added to: diuretics, antihypertensives, digitalis, converting-enzyme inhibitors, calcium channel blockers, beta-blockers,

Crite Lifus. Summy semants.

Crite Lifus. Summy semants.

Crite Lifus. Summy semants.

Criticoglycerin.

Endocrine Function: HMG-CoA reductase inhibitors interfere with cholesterol synthesis and lower circulating cholesterol levels and, as such, might theoretically blunt adrenal or gonadial steroid hormone production. Results of clinical trials with pravastatin in males and post-menopausal females were inconsistent with regard to possible effects of the drug on basal steroid hormone levels. In a study of 21 males, the mean testosterone response to human chorionic gonadotropin was significantly reduced (p<0.004) after 16 weeks of treatment with 40 mg of pravastatin. However, the percentage of patients showing a ≥50% rise in plasma testosterone after human chorionic gonadotropin stimulation did not change significantly after therapy in these patients. The effects of HMG-CoA reductase inhibitors on spermatogenesis and fertility have not been studied in adequate numbers of patients. The effects of HMG-CoA reductase inhibitors on spermatogenesis and fertility have not been studied in adequate numbers of patients. The effects of HMG-CoA reductase inhibitors on spermatogenesis and fertility have not been studied in adequate numbers of patients. The effects of HMG-CoA reductase inhibitors on spermatogenesis and fertility have not been studied in adequate numbers of patients. The effects of HMG-CoA reductase inhibitor on should be evaluated appropriately. Caution should also be exercised if an HMG-CoA reductase inhibitor or other agent used to tower cholesterol levels is administered to patients also receiving other drugs (e.g., ketoconazole, spironolactone, circultion) that may diminish the levels or activity of steroid hormones.

CNS Toxicity: CNS vascular lesions, characterized by perivascular hemorrhage and edema and mononuclear cell

infiltration of perivascular spaces, were seen in dogs treated with pravastatin at a dose of 25 mg/kg/day, a dose that produced a plasma drug level about 50 times higher than the mean drug level in humans taking 40 mg/day. Similar CNS vascular lesions have been observed with several other drugs in this class.

A chemically similar drug in this class produced optic nerve degeneration (Wallerian degeneration of retinogeniculate fibers) in clinically normal dogs in a dose-dependent fashion starting at 60 mg/kg/day, a dose that produced mean plasma drug levels about 30 times higher than the mean drug level in humans taking the highest recommended dose (as measured by total enzyme inhibitory activity). This same drug also produced vestibulocochiear Wallerian-like degeneration and retinal agnigion cell chromatolysis in dogs treated for 14 weeks at 180 mg/kg/day, a dose which resulted in a mean plasma drug level similar to that seen with the 60 mg/kg/dose.

Carcinogenesis, Mutagenesis, Impairment of Fertility: in a 2-year study in rats led pravastatin at doses of 10, 30, or 100 mg/kg body weight, there was an increased incidence of hepatocellular carcinomas in males at the highest dose (p<0.01). Although rats were given up to 125 times the human dose (HD) on a mg/kg body weight basis, their serum drug levels were only 6 to 10 times higher than those measured in humans given 40 mg pravastatin as measured by AUC.

The oral administration of 10, 30, or 100 mg/kg (producing plasma drug levels approximately 0.5 to 5.0 times human drug levels at 40 mg) of pravastatin to mice for 22 months resulted in a statistically significant increase in the incidence of malignant lymphomas in treated females when all treatment groups were pooled and compared to controls (p<0.05). The incidence was not dose-related and male mice were not affected.

A chemically similar drug in this class was administered to mice for 72 weeks at 25, 100, and 400 mg/kg body weight, which resulted in mean serum drug levels approximately 3, 15, and 33 times

creased spermatogenesis, spermatocytic degeneration, and giant cell formation in dogs. The clinical significance of these findings is unclear.

Pregnancy: Pregnancy: Category X: See CONTRAINDICATIONS.

Safety in pregnant women has not been established. Pravastatin was not teratogenic in rats at doses up to 1000 mg/kg daily. These doses resulted in 20x (rabbit) or 240x (rat) the human exposure based on surface area (mg/meter?). However, in studies with another HMG-COA reductase inhibitor, skeletal malformations were observed in rats and mice. PRAWACHOL (pravastatin sodium) should be administered to women of child-bearing potential only when such patients are highly unlikely to conceive and have been informed of the potential hazards. If the woman becomes pregnant while taking PRAWACHOL, it should be discontinued and the patient advised again as to the potential hazards to the fetus.

Nursing Mothers: A small amount of pravastatin is excreted in human breast milk. Because of the potential for serious adverse reactions in nursing infants, women taking PRAWACHOL, should not nurse (see CONTRAINDICATIONS).

Padiatric Use: Safety and effectiveness in individuals less than 18 years old have not been established. Hence, treatment in patients less than 18 years old is not recommended at this time. (See also PRECAUTIONS: General.) ADVERSE REACTIONS

ADVERSE REACTIONS

Pravastatin is generally well tolerated; adverse reactions have usually been mild and transient. In 4-month long placebo-controlled trials, 1.7% of pravastatin-treated patients and 1.2% of placebo-treated patients were discontinued from treatment because of adverse expenences attributed to study drug herapy, this difference was not statistically significant. In long-term studies, the most common reasons for discontinuation were asymptomatic serum transaminase increases and mild, non-specific gastrointestinal complaints. During clinical trials the overall incidence of adverse events in the eliderly was not different from the incidence observed in younger patients.

Adverse Clinical Events: All adverse clinical events (regardless of attribution) reported in more than 2% or pravastatin-treated patients in the placebo-controlled trials are identified in the table below, also shown are the percentages of patients in whom these medical events were believed to be related or possibly related to the drug:

Body System/Event	All Events %		Events Attributed to Study Drug %	
	Pravastatin (N = 900)	Placebo (N = 411)	Pravastatin (N = 900)	Placebo (N = 411)
Cardiovascular				
Cardiac Chest Pain	4.0	3.4	0.1	0.0
Dermatologic				
Rash	4.0°	1.1	1.3	0.9
Gastrointestinal				
Nausea/Vomiting	7.3	7.1	2.9	3.4
Diarrhea	6.2	5.6	2.0	1.9
Abdominal Pain	5.4	6.9	2.0	3.9
Constipation	4.0	7.1	2.4	5.1
Flatulence	3.3	3.6	2.7	3.4
Heartburn	2.9	1.9	2.0	0.7
General		-	=	=
Fatique	3.8	3.4	1.9	1.0
Chest Pain	3.7	1.9	0.3	0.2
Influenza	2.4*	0.7	0.0	0.0
Musculoskeletal		-		
Localized Pain	10.0	9.0	1.4	1.5
Myalgia	2.7	1.0	0.6	0.0
Nervous System		****		
Headache	6.2	3.9	1.7*	0.2
Dizziness	3.3	3.2	1.0	0.5
Renal/Genitourinary	0.0	3.2		5.0
Urinary Abnormality	2.4	2.9	0.7	1.2
Respiratory		2.0	0.,	
Common Cold	7.0	6.3	0.0	0.0
Rhinitis	4.0	4.1	0.1	0.0
Cough	2.6	1.7	0.1	0.0

*Statistically significantly different from placebo.

Colugn

Statistically significantly different from placebo.
The following effects have been reported with drugs in this class:
Skeletal: myopathy, rhabdomyolysis.

Neurological: dysfunction of certain cranial nerves (including alteration of taste, impairment of extra-ocular movement, facial paresis), termor, vertigo, memory loss, paresthesia, peripheral neuropathy, peripheral nerve palsy.

Hypersensitivity Reactions: An apparent hypersensitivity syndrome has been reported rarely which has included one or more of the following features: anaphylaxis, angioedema, lupus erythematous-like syndrome, polymyagid returnatics, vasculitis, purpura, thrombocytopenia, leutopyria, anemotyric anemia, positive ANA, ESP increase, arthritis, arthralgia, urticaria, asthenia, photosensitivity, feer, chilis, flushing, malaise, dyspnea, toxic epidermal necrolysis, erythema multiforme, including Stevens-Johnson syndrome.

Gastrointestinal: pancreatitis, hepatitis, including chronic active hepatitis, cholestatic jaundice, fatty change in liver, and, rarely, cirrhosis, fulminant hepatic necrosis, and hepatioma, anorexia, vomiting.

Reproductive: gynecomastia, loss of libido, erectile dysfunction.

Eye: progression of cataracts (lens opacities), ophthalmoplegia.

Laboratory Test Abnormalities: Increases in serum transaminase (ALT, AST) values and CPK have been observed (see WARNINGS).

Transient, asymptomatic eosinophilia has been reported. Eosinophi counts usually returned to normal despite continued therapy. Anema, thrombocytopenia, and leukopenia have been reported with other HMG-CoA reductase inhibitors achieved with lovastatin or pravastatin is not associated with greater reduction in LDL-cholesterol than that achieved with lovastatin or pravastatin is not associated with greater reduction in LDL-cholesterol than that achieved with lovastatin or pravastatin alone. No adverse reactions unique to the combination or in addition to those previously reported for each drug alone have been reported. Myopathy and rhabdomyolysis (with or wi immunosuppressive drugs, gemfibrozii, erythromycin, or lipid-lowering doses of nicotinic acid. Concemitant therapy with HMG-CoA reductase inhibitors and these agents is generally not recommended. (See WARNINGS: **Skeletal Muscle** and PRECAUTIONS: **Drug Interactions.**)

OVERDOSAGE

e have been no reports of overdoses with pravastating Should an accidental overdose occur, treat symptomatically and institute supportive measures as required

Effective lipid management doesn't have to be tough



- Improves key lipids significant reduction in LDL-C¹

- Excellent safety profile
 Easy for patients once-baily dosing, well tolerated
 Usual dose: 20 mg once daily at bedtime, with or without food

pravastatin socium 20 mg tablets

PRAVACHOL is indicated as an adjunct to diet for the reduction of elevated total and LDL-cholesterol levels in patients with primary hypercholesterolemia (Types IIa and librarhen the response to diet alone has not been adequate. Active liver disease or unexplained transaminase elevation incy and lactation are contraindications to the use of pravastatin sodium.

Please see CONTRAINDICATIONS, WARNINGS, PRECAUTIONS, and ADVERSE REACTIONS in the brief summary of prescribing information on the adjacent page.



Bristol-Myers Squibb Company